ECE440 - Midterm Review Notes

Probability review

You should be fuent with notions such as probability spaces, the axioms of probability and their consequences, the law of total probability and Bayes' rule, discrete and continuous random variables (RVs), expectations, independence, conditional probability and expectations, and notions of convergence of sequences of RVs.

Some noteworthy concepts and results you should make sure you are comfortable with are:

1) E and F are mutually exclusive (disjoint) events if $E \setminus F = ;$. From the axioms of probability this implies that for disjoint E and F, then $P[E [F] = P[E] + P[F]$.
2) E and F are independent events if $P[E \setminus F] = P[E]P[F]$.
3) You should be able to tell the difference between discrete and continuous RVs.
4) A Bernoulli RV X with parameter p indicates that a random event E occurred (a "success"), where $P[E] = p$. Recall the pmf, that $E[X] = p$ and $var[X] = p(1 - p)$.
5) A Geometric RV X with parameter p counts the number of independent Bernoulli trials till we register the frst success. Recall the pmf, that $E[X] = 1=p$ and $var[X] = (1 - p)=p^2$.
6) A Binomial RV X with parameters n and p counts the number of successes in n independent Bernoulli trials. Recall that $E[X] = np$ and $var[X] = np(1 - p)$. For $B_1; : : : ; B_n$ i.i.d. Bernoulli RVs with parameter p, one can write
$$X = \sum_{i=1}^{n} B_i :$$
7) A Poisson RV X with parameter $$ counts rare events or "arrivals". Recall the pmf, that $E[X] =$ and $var[X] =$. If X_1 Poisson($_1$) and X_2 Poisson($_2$) are independent, then $Y = X_1 + X_2$ Poisson($_1 +$ $_2$). The law of rare events asserts that the distribution of X Binomial$(n; p)$ converges to a Poisson$()$ as $n ! 1$, provided $np =$.
8) A Gaussian (Normal) RV X with parameters and ^{2} models randomness arising from the superposition of large number of random effects. (This statement should be clear if you understood the Central Limit Theorem.) Recall the pdf, that $E[X] =$ and $var[X] =$ ^{2}.
9) An indicator RV $I f E g$ indicates the occurrence of event E. Indicator RVs are Bernoulli distributed, with parameter $p = P[E]$. You should be fuent in operating with indicator RVs. The expectation of an indicator RV is the probability of the indicated event, i.e., $E[I f E g] = P[E]$.
10) It is useful to master the properties of expected values, such as the "law of the unconscious statistician" to calculate the expected value of a function of a RV, $E[f(X)]$. The expected value is a linear operator, hence for scalars a_i and b, $i = 1; : : : ; n$, then
$$E\left[\sum_{i=1}^{n} a_i X_i + b\right] = \sum_{i=1}^{n} a_i E[X_i] + b :$$
In general, expectation can be interchanged with any linear operation. If X and Y are independent RVs, then for any functions $f(\cdot)$ and $g(\cdot)$ one has
$$E[f(X)g(Y)] = E[f(X)]E[g(Y)] :$$

Expectation can be interchanged with products only if the RVs are independent, but it can always be interchanged with sums. Also, for an independent sequence of RVs $X_1, \ldots, X_n$ and scalars a_i and b_i, $i = 1, \ldots, n$, then

$$\text{var}\left[\sum_{i=1}^{n} a_i X_i + b\right] = \sum_{i=1}^{n} a_i^2 \text{var}[X_i].$$

11) The covariance is a measure of linear dependence between pairs of RVs, given by

$$\text{cov}[X, Y] = E[(X - E[X])(Y - E[Y])] = E[XY] - E[X]E[Y].$$

Recall that independent RVs are also uncorrelated, but the converse is not true in general (it is true though, if the random variables are jointly Gaussian).

12) Make sure you understand the statements of Markov's and Chebyshev's inequalities.

13) You should have an intuitive understanding of the difference between almost sure convergence, convergence in probability and convergence in distribution. Recall that the limit of a sequence of RVs is itself a RV.

14) The simplest class of random processes is i.i.d. sequences of RVs. You should understand and be able to apply the two main limiting theorems that pertain to sums of i.i.d. RVs. The Strong Law of Large Numbers asserts that for a sequence $X_N = X_1, X_2, \ldots$ of i.i.d. RVs with mean $E[X_1] = \mu$, the sample average converges to the mean almost surely, i.e.,

$$P\left[\lim_{n\to\infty} \frac{1}{n}\sum_{i=1}^{n} X_i = \mu\right] = 1.$$

If the i.i.d. RVs have variance $E[(X_1 - \mu)^2] = \sigma^2$, the Central Limit Theorem asserts that

$$\lim_{n\to\infty} P\left[\frac{\sum_{i=1}^{n} X_i - n\mu}{\sigma\sqrt{n}} \le x\right] = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x} e^{-u^2/2}\, du.$$

This implies the above normalized sum of RVs converges in distribution to a standard Gaussian RV.

15) Conditional probabilities and expectations and very useful tools for the analysis of random processes. Make sure you are comfortable with the defnitions and examples from the lecture slides, and that you understand the related problems from the homework assignments.

16) The conditional expectation $E[X|Y]$ is a function of Y, hence a RV which takes the values $E[X|Y = y]$. The law of iterated expectations states that $E[X] = E_Y[E_X[X|Y]]$, and is often helpful in computing expected values by conditioning. Likewise, one can compute the variance by using the formula

$$\text{var}[X] = E_Y[\text{var}_X[X|Y]] + \text{var}_Y[E_X[X|Y]].$$

Markov chains

You should be fuent with notions such as the Markov property, (multi-step) transition probabilities, representing a Markov chain (MC) through the transition probability matrix or state transition diagram, the Chapman-Kolmogorov equations, computing n-step transition probabilities and unconditional probabilities, classifying states of a MC, identifying communicating classes and irreducible MCs, computing stationary distributions, and understanding as well as applying